

ECTOKIL

**Water Soluble Ivermectin 1%
for External & Internal Parasites**

COMPOSITION :

Each 1000 gm. contains:

Ivermectin : 1% w/w
Livamisole hydrochloride : 1%
Lactose as carrier : upto 1000 gms.

INDICATIONS :

Useful for the treatment & control of infestations caused by internal and external parasites in Poultry & Veterinary.

DOSAGE : Poultry :

Through Drinking Water :

(0.4 mg of Ivermectin / kg. body weight)

Internal Parasitism :

Add 1 gm powder of "ECTOKIL" in drinking water required for each 25 kg. of body weight. If necessary repeat the treatment in 15 days.



External Parasitism :

Add 1 gm powder of "ECTOKIL" in drinking water required for each 25 kg. of body weight, for two consecutive days. If necessary repeat the treatment in 15 days.

Large Animals : 10 gm / 400 kg. body weight.
(or) as advised by a veterinarian.

PRECAUTIONS FOR USE : Medicated water must be administered within the next 12 hours.

Withdrawal Period : Broilers : 8 days

PRESENTATION : 100 gm.



Manufactured in India by :

**international
BIOLOGICALS**

Corp. Office : 40-9-60, Kohinoor Apts., Kalanagar,
Benz Circle, VIJAYAWADA - 520 010. Andhra Pradesh, INDIA.
Tel : +91 866 2483927, 6668855. www.intbio.in

