

ORGACAL-P PLUS

Calcium, Phosphorus with Vitamins

Composition: Each gram contains:

Elemental Calcium	: 45 mg
Elemental phosphorus	: 16 mg
Vitamin D ₃	: 400 IU
Vitamin B ₁	: 2.00 mg
Vitamin B ₂	: 2.00 mg
Vitamin B ₁₂	: 10 mcg
Vitamin C	: 6.00 mg
Citric acid	: 0.28 gm

Elemental calcium (45 mg): It is very important, structural, constituent of bones, it is essential for neuromuscular excitability, it controls cell permeability & it is co factor for many enzymes.

Elemental phosphorus (16 mg): It ensure skeletal growth. It is important mineral in bones. It increases calcium deposition, it prevents leg weakness, lameness etc.

Vitamin D₃ (400 IU): It plays an important role in the regulation of calcium and phosphorus metabolism.

Vitamin B₁ (2 mg): Vitamin B₁ plays a crucial role in the metabolism of carbohydrates and proteins to produce energy for the body. As well it helps to maintain nervous & cardiovascular system.

Vitamin B₂ (2 mg): Vitamin B₂ helps to convert carbohydrates into energy. It plays very important role in the production of red blood cells.

Vitamin B₁₂ (10 mg): It is essential vitamin, required for, hemoglobin & red blood cell formation, for hepatic regeneration & for general growth of the animal.

Vitamin C (6 mg): It is essential in the formation and maintenance of skeleton tissue and collagen. It participates as an oxidation reduction agent in numerous cellular process. It increases, the body's resistance to infections. The special application of vitamin C is to reduce stress during all kind of stress.

Citric acid (280 mg): It provides, acidic medium, for the control of pathogens in gut region.

Benefits:

■ Better utilization of calcium and phosphorus ■ Prevents leg weakness, lameness, osteomalacia, perosis, egg breakages, thin shelled eggs, rough shelled eggs ■ Increases growth, body weight and bone strength in broilers ■ Increases egg size, egg shell quality and egg production in layers ■ Increases hatchability in breeders ■ Prevents cage layer fatigue, prolapse, pecking, cannibalism etc.

Indications :

● Legs weakness, lameness ● Poor growth ● Egg breakages, rough shelled eggs, thin shelled eggs ● Low production ● Poor feathering ● Poor hatchability ● Pecking and cannibalism ● Cage layer fatigue ● Prolapse etc.


Dosage:

Poultry: 1 gm / 2 Ltrs of drinking water for 5-7 days

Livestock :

Milch Animals (Cow & Buffaloe) : 50 to 60 gm / animal, twice daily, **Calves, Sheep & Goat :** 20 to 30 gm / animal, twice daily, **Draught Animals :** 60 - 80 gm / animal, twice daily, **Dog :** 10-15 gm / animal, twice daily (or) as advised by a veterinarian (or) as advised by a veterinarian.

Presentation: 500gm, 1kg. plastic tin, 25 kg. bag



Manufactured by :



**international
BIOLOGICALS**

54-10-13(B), Phase-III, Telephone Exchange Road,
Jawahar Autonagar, Vijayawada-520 007.
Cell : +91 98481 56569, website : www.intbio.in

CERTIFICATIONS
cGMP
&
ISO 9001 : 2008



Marketed by:

Safe Bio Products Ltd
Mowna, Gazipur, Bangladesh