



TILMICODEM 25

Oral solution for poultry

- Excellent solubility
- Long lasting stability
- Proven efficacy in tylosin resistant pathogens



Oral solutions range

www.delosmedica.com

NAME OF THE VETERINARY**MEDICINAL PRODUCT:**

TILMICODEM 25, as tilmicosin phosphate
250 mg/ml, oral solution for poultry.

of Tilmicodem 25 should be calculated
according to the following formula:

$$\frac{\text{ml Tilmicodem 25 / kg body weight/day} \times \text{mean body weight of animals to be treated (kg)}}{\text{Mean daily water consumption (liters) / animal}} = \frac{\text{ml Tilmicodem 25}}{\text{liter of drinking water}}$$

STATEMENT OF THE**ACTIVE SUBSTANCE****AND OTHER INGREDIENTS:**

1 ml solution contains:

Tilmicosin phosphate: 250 mg (equivalent to 225 mg tilmicosin base), Excipients (propyl gallate, disodium edetate, orthophosphoric acid, purified water) ad. 1 ml.

ADVICE ON CORRECT**ADMINISTRATION:**

The product must be diluted before administration to the animals. During treatment, the animals should drink only medicated water. Medicated drinking water should be prepared fresh every 24 hours.

INDICATIONS

In poultry: for treatment of infections produced by susceptible bacteria such as: *Mycoplasma* spp, *Pasteurella multocida*, *Haemophilus paragallinarum*, *Streptococcus* spp., anaerobes (*Clostridium* spp.) etc.

WITHDRAWAL PERIOD:

Edible tissues of poultry: 13 days from cessation of oral medication. Not permitted for use in laying bird producing eggs for human consumption.

CONTRAINDICATIONS:

Do not allow horses and other equines access to drinking water containing Tilmicodem 25.

SPECIAL PRECAUTIONS**FOR STORAGE:**

Keep away from reach and sight of children! Store in original packaging, at room temperature (below 25°C), protected from moisture and direct sunlight. Shelf life of the veterinary medicinal product as packaged for sale: 2 years. Shelf life after dilution in drinking water, according to directions: 24 hours. Shelf-life after first opening of package: 28 days. Close tight the recipient after use. Do not use after expiry date which is stated on the label.

TARGET SPECIES:

Poultry.

DOSAGE FOR EACH SPECIES,**ROUTE(S) AND METHOD****OF ADMINISTRATION**

In poultry: the product is administered via drinking water in dosage of 15 - 20 mg tilmicosin/kg (equivalent to 17 - 22 mg tilmicosin phosphate/kg b.w.) corresponding to 0.07 - 0.09 ml Tilmicodem 25/kg b.w./day for 3 - 5 consecutive days. Based on the recommended doses and the number and weight of the animals to be treated, the exact daily amount