



VERMICID 10

Oral suspension for cattle, sheep and Poultry

- Proper concentration
- Proven efficacy
- 100% compliance with E.Ph. 6



Oral solutions range

www.delosmedica.com

NAME OF THE VETERINARY MEDICINAL PRODUCT

VERMICID 10, 100 mg/g, oral suspension for Cattle, Sheep and Poultry. Albendazole.

STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS PER ML:

Albendazole 100 mg,

Excipients: Benzoic acid 8 mg

INDICATIONS:

The product Vermicid 10 is indicated for the control of albendazole susceptible mature and developing immature forms of the following internal parasites of cattle, sheep and Poultry.:

DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION:

To ensure administration of a correct dose, body weight should be determined as accurately as possible; accuracy of the dosing device should be checked. The therapeutically dosage is approximately 10 – 12.5 mg albendazole per kg bodyweight.

Cattle: 5 ml / 40 kg b.w. (12.5 mg albendazole/kg bw); Sheep: 4 ml/40 kg b.w. (10 mg albendazole/kg bw); Poultry : 1 ml/10 kg b.w.(10 mg albendazole/kg bw).

WITHDRAWAL PERIOD:

Cattle: 28 days for meat; Sheep: 4 days for meat; Milk: 3 days, chicken/ egg: 7 days

Parasitosis	Cattle	Sheep	Poultry
Control of adult liver flukes	Fasciola hepatica, dicrocoelium, Lancelatum	Fasciola hepatica, dicrocoelium, Lancelatum	
Control of tapeworms	Moniezia benedeni, M. expansa	Moniezia expansa	Raillitena Sp.
Gastro-intestinal roundworms(L4 larvae and adults)	Ostertagia ostertagi(including inhibited larvae L4), Haemonchus contortus, Trichostrongylus axei	Ostertagia circumcincta(including inhibited larvae L4), Cooperia oncophora, Haemonchus contortus, Trichostrongylus axei	Ascaris Sp. Capillaria Sp. Heterakis gallinarum,
Gastro-intestinal roundworms(L4 larvae and adults)	Nematodirus spathiger, Cooperia oncophora(including inhibited larvae L4)	Nematodirus spathiger, N. Filifolis, cooperia oncophora, Trichostrongylus colubriformis, Oesophagostomum, Columbianum, Chabertia ovina	Ascaris Sp. Capillaria Sp. Heterakis gallinarum
Gastro-intestinal adult roundworms	Bonostomum phlebotomum, Trichostrongylus colubriformis, Oesophagostomum radiatum.		Ascaris Sp. Capillaria Sp. Heterakis gallinarum
Lungworms(Larva & adults)	Dictiocaulus viviparus	Dictiocaulus filaria	Syngamas trachea

The product has also ovicidal activity. As prophylactic purposes, the deworming program should consist in 3-4 administrations: before entering to pasture, in July and September and after housing the animals.

CONTRAINDICATIONS:

Do not administer to animals with known hypersensitivity to benzimidazoles or the excipients.

TARGET SPECIES:

Cattle, Sheep and Poultry.

Animals must not be slaughtered for human consumption during treatment. Milk for human consumption must not be taken during treatment.

SPECIAL PRECAUTIONS FOR STORAGE:

Store in original packaging, at temperature between 15- 25⁰C, protected from direct sunlight and freeze. Keep away from reach and sight of children!

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.